

A decorative graphic featuring a central white scrollwork element on the left, a light blue flower at the top, and a red flower at the bottom, all surrounding a central white-bordered frame. The background is a repeating pattern of light orange roses.

What We Know

NEW SCIENCE LINKING
OUR HEALTH AND
THE ENVIRONMENT

what we know

NEW SCIENCE LINKING OUR HEALTH AND THE ENVIRONMENT

Helpful Definitions

ENVIRONMENTAL CONTAMINANTS

A broad range of substances that may harm people's health, including synthetic chemicals and chemical compounds, heavy metals (used in products or industrial processes) and radiation. The term also includes air pollution, which is a mixture of particles and chemicals from different sources, including industrial pollution, vehicle emissions (such as cars, diesel trucks and buses, freight ships and others) and tobacco smoke. Sometimes referred to as simply "contaminants" in this document.

CHEMICALS

More than 82,000 synthetic chemicals are registered for use in manufacturing and industry. Only ten percent of those chemicals have been tested for their effects on human health. Approximately 2,800 of these are considered "high-production volume" chemicals, or those produced in amounts of 1 million pounds or more per year in the United States.

Women play many roles. We are friends, sisters, daughters, girlfriends, partners, wives, workers and mothers. We are caretakers and breadwinners. We are advocates for our families, children, aging parents and communities.

Women also are leaders in protecting people's health—whether it is taking a sick child's temperature, making sure there is nutritious food in the house or working toward a healthy environment.

This booklet explains the science linking women's health to the environment. It describes the factors that affect whether contaminants will harm our health. It also highlights the effects of contaminants at each stage of a woman's life—including the potential life-long health effects from being exposed to contaminants in the womb.

The information presented here can be scary. There is reason to be concerned, but there are also many opportunities to improve the world we live in. We can use safer products around our homes, and we can hold government and industry accountable for protecting us from environmental contaminants. The other booklets in this series, *What You Can Do: Everyday Actions to Protect Your Health* and *What We Can Do: Community Efforts to Protect Our Health* provide ideas for taking action. All three booklets are available at www.womenshealthandenvironment.org/toolkit.



Environmental Contaminants and Our Health

Health problems and disabilities such as asthma, birth defects, cancer, diabetes and infertility affect more than 100 million men, women and children, about one-third of the U.S. population. Scientific research shows that exposure to environmental contaminants may increase a person's risk for these and other health concerns. Pregnancy and early childhood are especially vulnerable times, but exposure throughout our lives may harm us.

We can be exposed to these contaminants through household products, the food we eat, air pollution from vehicles or factories, hazardous waste sites and other sources. We can find contaminants in our homes, communities, schools and workplaces. With thousands of chemicals and other contaminants in use, more research is needed to understand how all these environmental contaminants affect human health. But there is enough evidence to know we need better public policy to prevent exposing people to contaminants that are—or may be—harmful to people's health.

If the annual average concentration of fine particulate air pollution could be reduced, thousands of premature deaths, heart attacks, strokes and bypass surgeries across the country could be prevented.¹



Pollution in People

One way scientists are getting a better understanding of how environmental contaminants are impacting people is through biomonitoring or “body burden” studies. These studies include testing people's urine, breast milk, blood and other biological samples to see if they contain contaminants.

Although biomonitoring cannot explain how people were exposed to contaminants or whether these exposures will lead to any health problems, the test results document the number and volume of contaminants that end up in our bodies. Some body burden findings include:

Lead in children: For the past three decades, the Centers for Disease Control and Prevention (CDC) have conducted body burden studies of lead, which may cause learning disabilities and other problems. The most recent studies found that lead levels in children have declined, indicating that efforts to reduce childhood exposure to lead in gasoline and other products have had success.² More work is needed, especially to protect children who live in homes contaminated with lead from paint or other sources.

Flame retardants in breast milk: A biomonitoring study by the Environmental Working Group (EWG) found that women in the United States have the world's highest levels of flame retardants in their breast milk³ Flame retardants, which are widely used in furniture, electronic equipment and other consumer products, may be associated with learning and memory problems and behavior changes. Several states have banned the use of certain flame retardants based on these biomonitoring studies.

Contaminants in newborns: A study by EWG and Commonwealth of ten newborns' umbilical cord blood showed an average of 200 industrial chemicals and pollutants. Overall, the study detected 287 out of the 413 chemicals that were surveyed.⁴ This result showed that not even the mother's placenta can keep environmental contaminants from crossing into her child.

Some Women's Health Concerns That May Be Linked to Environmental Contaminants

There is growing evidence that some of the health concerns that especially affect women may be linked to environmental contaminants. Some examples include:

Breast Cancer: A woman has a one in eight chance of having breast cancer in her lifetime,⁵ but less than one in 10 of those women are born with a genetic predisposition for the disease.⁶ Radiation (including medical radiation)^{7 8} and longer lifetime exposures to the hormone estrogen, and perhaps also to progesterone,⁹ can increase a woman's risk for breast cancer. Industrial chemicals that act like the hormones estrogen and progesterone may be adding to women's breast cancer risk. Other factors may also play a role.¹⁰

Heart Disease: Cardiovascular disease remains the leading cause of death in the United States, and the No. 1 cause of death for women.¹¹ Some of the environmental factors that increase the risk for heart disease include smoking¹² and secondhand smoke,¹³ exposure to some metals (such as arsenic,^{14 15 16 17} lead^{18 19} or mercury^{20 21 22}) and air pollution.^{23 24 25}

Endometriosis: Endometriosis affects about 1 in 10 women in the United States,²⁶ and nearly 40 percent of women with infertility appear to have endometriosis.²⁷ In this disease, the inner lining of the uterus, called the endometrium, is found outside the uterus in the abdominal cavity or on the ovaries or fallopian tubes. Endometriosis may cause pain during menstruation or intercourse. Some contaminants that may increase risk for endometriosis include dioxins,^{28 29 30} polychlorinated biphenyls (PCBs)^{31 32} and radiation.^{33 34}

Fertility Challenges: According to a recent national survey, 12 percent of the reproductive age population in the United States, or 7.3 million couples, report difficulty conceiving or carrying a pregnancy to term, or both of these problems. The largest reported increase in infertility is in women under age 25.³⁵ Although there is still much to learn about environmental factors, exposures that may interfere with fertility include cigarette smoke,^{36 37} DES exposure in the womb (some women took the drug DES to prevent miscarriages from the 1930s to 1971)³⁸ and health challenges such as endometriosis.³⁹ Exposure to solvents such as formaldehyde,⁴⁰ toluene,^{41 42 43} perchloroethylene⁴⁴ (dry cleaning fluid) and pesticides⁴⁵ also has been linked to fertility problems in women.

To learn more about which environmental contaminants may increase risk for these and other health conditions, see the Collaborative on Health and the Environment's Toxicant and Disease database:
<http://database.healthandenvironment.org>.



Diabetes: There has been a 46 percent increase in women with diabetes since the mid-1990s.⁴⁶ Nutrition and obesity are significant risk factors for diabetes.⁴⁷ Research is showing that environmental contaminants may also increase risk for diabetes. For example, exposure to estrogen-like compounds is associated with the development of obesity, therefore increasing risk of developing diabetes.⁴⁸ Arsenic, and to a lesser extent dioxin, have both been associated with higher rates of adult onset of Type II diabetes.^{49 50 51} A recent animal study showed that chronic exposure to low levels of Bisphenol A (found in polycarbonate plastic and food can linings) causes insulin resistance in adult mice, a phenomenon linked to Type II diabetes.⁵² Furthermore, new studies are finding that exposure to PCBs and organochlorine pesticides may also increase risk for diabetes.⁵³

Health Disparities Due to Race and Ethnicity

Health concerns and disease can affect all people, but they do not affect all people equally. For example, children of color suffer disproportionate burdens of disease with potential environmental aspects, including asthma, learning and developmental disabilities and cancer.⁵⁴ African American women are more likely to die from heart disease, stroke or cancer than women of any other race.⁵⁵

Health Factors: A Complex Web

A woman's health is a balance of many complex factors. Exposure to environmental contaminants is just one of them. There is no formula that can determine the exact effects of exposure to different contaminants. The effect depends on the risk associated with that exposure and a person's overall health and susceptibility. This illustration shows just some of the factors at work.

To protect our health, it is important to do the positive things where we have control, such as eating well, exercising, using healthier products and reducing stress. And then join the growing movement of women working to make sure the government does its job of protecting all of us from environmental contaminants.



Environmental Contaminants

It is impossible to make exact predictions of whether a person exposed to environmental contaminants will develop health problems. But some of the factors that influence the risk of being harmed by contaminants include:

Hazard of the Contaminant

Not all contaminants are equally hazardous. Adequate information about the health impacts of thousands of chemicals does not yet exist; yet it is clear that some contaminants are especially hazardous to human health and the health of other species. For example, mercury is widely understood to affect brain development, especially in the very young.

Dose of Exposure

For many years people assumed that contaminants were only harmful if people were exposed in high doses. But new research indicates that the question of dose is more complicated. Some contaminants are harmful at high doses, some at low doses, some at both high and low doses, but in different ways.

Timing of Exposure

Exposures to some contaminants early in life can influence health later in life. For example, early life exposure to chemicals that are hormonally-active (i.e. interfere with the hormone system), such as Bisphenol A, may cause puberty to happen earlier in life.^{56 57 58 59} Exposure to arsenic in the womb is tied to lung disease and cancer in adults.⁶⁰

Combinations of Exposures

Most of the research on the connection between health and contaminants is focused on exposure to one contaminant at a time. In reality, people are generally exposed to multiple contaminants at the same time through the variety of products we use, the food we eat, the emissions we are surrounded by and the contaminants that may be stored in our bodies for years.

Much more research is needed on how combinations of contaminants can affect our health, but smoking is one area where this kind of research has been done. The results are not good. For example, smokers exposed to asbestos have a markedly increased risk for lung cancer, much larger than the risk of lung cancer from just smoking or just asbestos exposure.⁶¹



Nutrition

Good nutrition is important for children to grow strong and for all people to support their overall health. Studies indicate that poor nutrition may magnify the effect of environmental contaminants. For example, iron-deficient children may experience greater effects from lead exposure because they actually absorb and retain more lead. Additionally, iron-deficient kids may ingest more lead because they are more likely to live in lead-contaminated environments. When, like any normal child, they put their hands in their mouths, they are likely to swallow more lead-contaminated dust and dirt.⁶²



Stress

Stress comes in many forms, such as too many responsibilities, financial pressure or challenges in our relationships. Reducing stress is important for our overall health and being able to handle life's challenges—mental, physical or otherwise. Constant stress taxes the immune system and makes us more vulnerable to harmful influences on our health, such as environmental contaminants. Taking time to exercise, eat well and do things that make you happy can help reduce stress.



Pregnancy

Events in a woman's reproductive cycle can affect her health. For example, women who have their first child after age 30 and women who never give birth have about a two- to three-fold increased risk of breast cancer compared to women who have their first child before age 20.⁶³



Exercise

Exercise is important for staying healthy. It builds strength, increases circulation and strengthens the immune system—all important things when fighting the everyday cold or resisting the effects of environmental contaminants. Plus it feels good and helps lift the spirits. However, care should be taken when exercising outdoors on poor air-quality days.^{64 65}



Social Support

Strong social support from friends, family and/or a supportive community can play an important role in a person's well-being and even help protect people against chronic illness.⁶⁷ For example, people with a strong social network may be less likely to develop obesity and diabetes.⁶⁸ One study found that the social support African American women received in their faith-based community helped them maintain healthier lifestyles.⁶⁹



Genetic Heritage

Each of us inherits our genes from our biological parents. Some of these genes determine how we look, such as eye color or face structure. Other genes make people more susceptible to developing some diseases. For example, people with inherited alterations in the genes called BRCA1 and BRCA2 (short for breast cancer 1 and breast cancer 2) have an increased risk for breast and ovarian cancer.⁶⁶ But carrying these genes does not mean a woman will automatically develop such cancers.

Each stage of a woman's life sets the path for her future health and possibly the health of her children. The previous section highlights the many factors that influence whether or not exposure to environmental contaminants will contribute to health problems. This section highlights how the timing of exposure may affect our health and the health of future generations.

Lifecycle

and the Environment



In Utero

Before a baby is born, critical development happens in all organs and tissues, including the rapidly growing brain, reproductive and endocrine (or hormone) systems. What babies are exposed to in the womb may have long-term health effects.

- The risk for attention deficit disorder and learning disabilities can be increased if a fetus is exposed to certain contaminants.⁷⁰ For example, there are some fish that pregnant women should avoid because they contain methylmercury,⁷¹ which can adversely affect brain development.^{72 73 74 75} But there are types of fish that pregnant women should eat because they are rich in important nutrients.
- Exposure to indoor and outdoor air pollutants (including secondhand smoke) has been linked to adverse birth outcomes, such as low birth weight and increased risk for developing asthma.^{76 77 78 79 80 81 82 83}
- Exposure to chemicals in the womb that affect the endocrine system may increase the risk of numerous types of cancer later in life.⁸⁴ For example, animal studies indicate that pre-natal exposure to Bisphe-nol A (which is used in polycarbonate plastics, the lining of food cans and other products) may increase the risk of breast cancer later in life.^{85 86}



Infants & Children

More growth and development takes place in a baby's first year than at any other time. Early childhood is also an important period of development. The early years are a critical time to provide good nutrition and a healthy environment.

- Chemicals in secondhand smoke are very damaging to a young child's health. They have been linked to increased risk for sudden infant death syndrome, middle ear infections, lower respiratory tract illness, prevalence of wheeze and cough and worsening asthma.^{87 88 89 90}
- Once children can crawl, they are exposed to floor dust that may contain lead, flame retardants or other substances that may harm brain development.^{91 92 93 94 95}
- When it comes to eating, children are not just little adults. Health impacts from food contaminants such as pesticides or heavy metals can be greater because children consume more food per pound of body weight than adults and because childhood is such a vulnerable time of development.⁹⁶
- Children may also be exposed to pesticides and harmful cleaners at school.



Teens

Teenage years are also a time of rapid growth. Girls' bodies begin producing higher levels of estrogen and progesterone, their reproductive tract undergoes change to prepare for pregnancy and their breasts go through rapid development. All of these changes make their bodies vulnerable to environmental contaminants.

- Teens often begin using makeup and personal care products, many of which contain ingredients that may be harmful to their health.^{97 98 99} While the chemicals present in any one cosmetic product alone may not cause harm, teenagers, like adults, use on average more than 10 personal-care products every day.
- Each day, nearly 6,000 children under 18 years of age start smoking, and 2,000 of them will become regular smokers. Approximately 90 percent of smokers begin smoking before age 21.¹⁰⁰ Studies indicate that when multiple factors are taken into account, such as the length of time a person has been a smoker and the amount she or he smokes, smoking during adolescence causes more damage than at other times, especially increasing the risk for lung cancer.¹⁰¹ Secondhand smoke may also increase risk for breast cancer, and developing breast tissue may be quite sensitive to environmental contaminants.¹⁰²



Early Adulthood

This is a time of adjusting to full-time jobs or other adult responsibilities. Finding ways to maintain good health habits while making the transition into adulthood is important, including eating well, maintaining or establishing an exercise routine and choosing healthier products around the home.

- Entering the workforce may entail being exposed to contaminants from toxic cleaners, chemicals used in manufacturing or other sources. Federal standards for occupational safety are often not strong enough and are not always well enforced. Workplace exposures may increase a woman's risk for many health concerns, such as cancer.¹⁰³
- Young women may be thinking of becoming a mother at some point in their lives, but exposures to chemicals even years before they try to have children may impact their ability to become pregnant and/or have a healthy, full-term child. For example, endometriosis, which may be caused by chemical exposures, can impair fertility.¹⁰⁴
105 106 107 108 109 110 111 112 Occupational exposure to certain chemicals is also known to threaten a woman's ability to have a healthy child.^{113 114 115 116}

Pregnancy

In the words of Katsi Cook, a Native American midwife, a woman's body is the first environment. Whatever a pregnant woman is exposed to, her baby may also be exposed to; the placenta does not protect the fetus from environmental contaminants.^{117 118} Eating nutritious food, exercising and avoiding environmental contaminants when possible are especially important during pregnancy.¹¹⁹

- Exposure to chemicals on the job when pregnant can be especially dangerous. For example, beauty salon workers and women working at dry cleaners may have a higher rate of miscarriage from contaminant exposures in the work place.^{120 121 122}
- Choosing healthier products around the house is especially important when a woman is pregnant. For example, studies have found that phthalates—chemicals used in vinyl plastic and some personal care products such as nail polish—may harm male babies' reproductive organs.¹²³ Pregnant women also should avoid using indoor pesticides, which can increase the baby's risk for being born prematurely and harm its development in the womb.

Menopause

A woman's changing hormones during menopause can cause obvious effects like hot flashes and night sweats, but a lifetime of exposure to contaminants may affect the onset of menopause and have other affects on a woman's health.

- A woman's risk for cardiovascular disease, including coronary heart disease and stroke, may increase as a result of menopause because estrogen, which decreases during menopause, serves as protection from these problems.^{124 125}
126 When this natural protection decreases over time, lifetime risk factors for cardiovascular disease such as exposure to environmental contaminants may be more pronounced.
- Menopause often accelerates bone disintegration which can release contaminants stored in the bone. For example, the release of lead stored in a woman's bones may impact her memory and her ability to think clearly.¹²⁷
- Chemical exposures throughout life may induce early or premature menopause and increase the risks of illness and death in the post-menopausal period.¹²⁸

Seniors

The likelihood of having a stroke, heart disease, cancer or most other illnesses, whether related to environmental contaminants or not, increases after age 65. The way a woman has treated her body her whole life and the environmental contaminants she has been exposed to can affect her health and well-being in her later years.

- It is normal for people's immune systems to grow weaker as they age,¹²⁹ but a lifetime of exposure to environmental contaminants may further suppress an older woman's immune system.¹³⁰
- Many women become less active as they age, which can lead to spending more time indoors and increase the chance of being exposed to indoor air pollutants. Effects from smoking or using cleaners and pesticides can be magnified if most of a woman's time is spent indoors in a poorly ventilated room.¹³¹
- Exposure to outdoor air pollution can also pose significant risk to older adults. For example, particle pollution and ozone (from car and other industrial emissions) may aggravate lung diseases.¹³²

What We Can Learn from Animals

In the 1960s, marine biologist and zoologist Rachel Carson changed the way Americans think about and protect the environment. By studying the effects of the pesticide DDT on animal populations, she warned us of the potential effects on human health. She broke new ground in 1962 when she published *Silent Spring*, which documented how birds' eggs were thinning due to DDT exposure, resulting in an alarming decrease in the bird's ability to reproduce. Her account of health concerns associated with pesticide use was one of the catalyzing moments for the environmental movement in the United States.

More than 40 years later, there is still plenty we can learn about environmental contaminants by monitoring animal populations in the wild. While what affects animals will not necessarily affect humans, the effects of contaminants on animal health certainly raises important concerns.



Frogs: Researchers have found evidence that atrazine—the most commonly used herbicide in the United States and probably in the world—can cause hermaphroditism (having both male and female reproductive organs) in frogs in the laboratory and may be responsible for the same impacts seen in frogs in the wild.^{133 134 135} Atrazine is widely used on corn and soybean fields in the United States and contaminates drinking water supplies.



Alligators: Studies of alligators in a Florida lake contaminated with DDT and other organochlorine pesticides in the 1980s found the males' penises were on average 25 percent smaller than normal. These alligators, at the top of the lake's food chain, ate fish that store the breakdown products of DDT. These byproducts accumulated in the alligators' bodies and caused changes in the hormonal system.^{136 137 138 139}



Birds: Hundreds of bird deaths have been reported from toxic fumes produced from normal household use of Teflon pans and products.¹⁴⁰ According to the Environmental Protection Agency, perfluorooctanoic acid (PFOA), one of the key compounds used in Teflon, is very persistent in the environment and can be found in the blood of the general U.S. population. Studies indicated that PFOA causes developmental and other adverse effects in laboratory animals.¹⁴¹



Polar Bears: Studies indicate that polar bears are threatened by chemicals accumulating in the environment. Initial studies indicate that polychlorinated biphenyls (PCBs) and other chlorine-based chemicals may be causing polar bears' ovaries and testicles to shrink,¹⁴² which may interfere with their ability to reproduce.

Learning about all the ways that environmental contaminants can impact health can be a little overwhelming. But there's good news. First, the human body is amazingly resilient. Doing your best to stay healthy through eating nutritious food and getting exercise helps your body stay strong. Second, as the other booklets in this series will show, there are proactive things we can do as individuals and as a community to reduce our exposures to environmental contaminants and improve the health of people and the planet we share.

For further details, go to www.womenshealthandenvironment.org.



Endnotes

- ¹ From a press release associated with the publication of: Miller KA, Siscovick DS, Sheppard L, Shepherd K, Sullivan JH, Anderson GL, Kaufman JD. *Long-Term Exposure to Air Pollution and Incidence of Cardiovascular Events in Women*. *New England Journal of Medicine* 2007 Feb 1; 356(5): 447-58.
- ² Centers for Disease Control and Prevention. *Third National Report on Human Exposure to Environmental Chemicals*. Atlanta (GA): CDC, 2005. www.cdc.gov/exposurereport/report.htm.
- ³ Environmental Working Group. *Mother's Milk: Record levels of toxic fire retardants in American mother's breast milk*. 2003. www.ewg.org/reports/mothersmilk/es.php. Viewed April 9, 2007.
- ⁴ Environmental Working Group. *Body Burden—The Pollution in Newborns*. 2005. www.ewg.org/reports/bodyburden2/execsumm.php. Viewed April 9, 2007.
- ⁵ American Cancer Society. *Breast Cancer Facts & Figures 2005-2006*. 2006.
- ⁶ Sasco A. *Epidemiology of breast cancer: An environmental disease?* *APMIS* 2001 May; 109(5):321-32.
- ⁷ U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. *Report on Carcinogens, Eleventh Edition*. 2005.
- ⁸ National Research Council. *Biologic effects of ionizing radiation VII: Health risks from exposure to low levels of ionizing radiation*. National Academy of Science, Washington DC. 2005.
- ⁹ Davis DL, Axelrod D, Osborne MP, Telang NT. *Environmental influences on breast cancer risk*. *Science and Medicine* 1997; 4:56-63.
- ¹⁰ See Vassar College Breast Cancer and the Environment Project. <http://erbc.vassar.edu>.
- ¹¹ American Heart Association. *Women, Heart Disease and Stroke*. 2007. www.americanheart.org/presenter.jhtml?identifier=4786. Viewed April 9, 2007.
- ¹² Centers for Disease Control and Prevention. *2001 Surgeon General's Report - Women and Smoking*. 2001. www.cdc.gov/tobacco/data_statistics/sgr/sgr_2001/index.htm.
- ¹³ Center for Disease Control and Prevention. *Secondhand Smoke Causes Heart Disease*. 2006. www.cdc.gov/tobacco/data_statistics/Factsheets/HeartDisease.htm. Viewed April 9, 2007.
- ¹⁴ Tseng W. *Effects and dose-response relationship of skin cancer and black foot disease with arsenic*. *Environmental Health Perspectives* 1977; 19:109-119.
- ¹⁵ Tseng C, Chong C, Tseng CP, Hsueh Y, Chiou H, et al. *Long-term arsenic exposure and ischemic heart disease in arseniasis-hyperendemic villages in Taiwan*. *Toxicology Letters* 2003 Jan 31; 137(1-2):15-21.
- ¹⁶ Rosenberg H. *Systemic arterial disease and chronic arsenicism in infants*. *Archives of Pathology* 1974; 97:360-365.
- ¹⁷ Zierold K, Knobloch L, Anderson H. *Prevalence of chronic diseases in adults exposed to arsenic-contaminated drinking water*. *American Journal of Public Health* 2004; 94(11): 1936-1937.
- ¹⁸ Cheng Y, Schwartz J, Sparrow D, Aro A, Weiss ST, Hu H. *Bone lead and blood lead levels in relation to baseline blood pressure and the prospective development of hypertension: the Normative Aging Study*. *American Journal of Epidemiology* 2001; 153(2):164-71.
- ¹⁹ Menke A, Muntner P, Batuman V, Silbergeld EK, Guallar E. *Blood lead below 0.48 micromol/L (10 microg/dL) and mortality among US adults*. *Circulation* 2006 Sep 26; 114(13):1388-94.
- ²⁰ Salonen JT, Seppanen K, Nyyssonen K, Korpela H, et al. *Intake of mercury from fish, lipid peroxidation, and the risk of myocardial infarction and coronary, cardiovascular, and any death in eastern Finnish men*. *Circulation* 1995; 91(3):645-55.
- ²¹ Virtanen JK, Voutilainen S, Rissanen TH, Mursu J, et al. *Mercury, fish oils, and risk of acute coronary events and cardiovascular disease, coronary heart disease, and all-cause mortality in men in eastern Finland*. *Arteriosclerosis Thrombosis Vascular Biology* 2005 Jan; 25(1):228-33.
- ²² Guallar E, Sanz-Gallardo MI, van't Veer P, Bode P, et al. *Mercury, fish oils, and the risk of myocardial infarction*. *New England Journal of Medicine* 2002; 347(22):1747-54.
- ²³ Brook R, Franklin B, Cascio WE, Hong Y, et al. *Air Pollution and Cardiovascular Disease. A Statement for Healthcare Professionals from the Expert Panel on Population and Prevention Science of the American Heart Association*. *Circulation* 2004; 109:2655-2671, 2004.
- ²⁴ Dockery DW, Pope CA 3rd, Xu X, Spengler JD, et al. *An association between air pollution and mortality in six U.S. cities*. *New England Journal of Medicine* 1993; 329(24):1753-9.
- ²⁵ Miller KA, Siscovick DS, Sheppard L, Shepherd K, Sullivan JH, Anderson GL, Kaufman JD. *Long-Term Exposure to Air Pollution and Incidence of Cardiovascular Events in Women*. *New England Journal of Medicine* 2007 Feb 1; 356(5): 447-58.
- ²⁶ Eskenazi B and M Warner. *Epidemiology of endometriosis*. *Obstetrics Gynecology Clinics of North America* 1997; 24(2):235-258.
- ²⁷ *Challenged conceptions: Environmental Chemicals and Infertility*. 2005. Collaborative on Health and the Environment.
- ²⁸ Mayani A, Barel S, Soback S, et al. *Dioxin concentrations in women with endometriosis*. *Human Reproduction* 1997; 12:373-375.
- ²⁹ Birnbaum LS, Cummings A. *Dioxins and endometriosis: a plausible hypothesis*. *Environmental Health Perspectives* 2002; 110(1):15-21
- ³⁰ Rier, S. and W.G. Foster, *Environmental dioxins and endometriosis*. *Seminars in Reproductive Medicine* 2003; 21(2): 145-54.
- ³¹ Rier S, Turner W, Martin D, et al. *Serum levels of TCDD and dioxin-like chemicals in Rhesus monkeys chronically exposed to dioxin: correlation of increased serum PCB levels with endometriosis*. *Toxicological Sciences* 2001; 59:147-159.
- ³² Louis GM, Weiner JM, Whitcomb BW, Sperrazza R, Schisterman EF, Lobbell DT, Crickard K, Greizerstein H, and Kostyniak PJ. *Environmental PCB exposure and risk of endometriosis*. *Human Reproduction* 2005 Jan; 20(1):279-85.
- ³³ Fanton J, Golden J. *Radiation-induced endometriosis in Macaca mulatta*. *Radiation Research* 1991; 126: 141-146.
- ³⁴ Palumbo G, Variiale L, Paba V, et al. *Effect of space radiation on expression of apoptosis-related genes in endometrial cells: a preliminary study*. *Physica Medica* 2001; 17 Suppl 1:241-6.
- ³⁵ Center for Disease Control. National Center for Health Statistics. *Fertility, Family Planning, and Reproductive Health of U.S. Women: Data from the 2002 National Survey of Family Growth*. 2002. www.cdc.gov/nchs/data/series/sr_23/sr23_025.pdf.
- ³⁶ Augood C, Duckitt K, Templeton A. *Smoking and female infertility: a systematic review and meta-analysis*. *Human Reproduction* 1998; 13:1532-1539.
- ³⁷ Olsen J, Rachtoun P, Schiødt A, et al. *Tobacco use, alcohol consumption, and infertility*. *International Journal of Epidemiology* 1983; 12:179-184.
- ³⁸ Giusti R, Iwamoto K, Hatch E. *Diethylstilbestrol revisited: a review of the long term health effects*. *Annals of Internal Medicine* 1995; 122:778-788.
- ³⁹ For more reading, see The Collaborative on Health and the Environment's infertility resource page. www.healthandenvironment.org/infertility/val-lombrosa_documents.

- ⁴⁰ Taskinen H, Kyronen P, Sallmen M, et al. *Reduced fertility among female wood workers exposed to formaldehyde*. American Journal of Industrial Medicine 1999; 36(1):206-212.
- ⁴¹ Ng T, Foo S, Yoong T. *Risk of spontaneous abortion in workers exposed to toluene*. British Journal of Industrial Medicine 1992; 49:804-808.
- ⁴² Plenge-Bonig A, Karmus W. *Exposure to toluene in the printing industry is associated with subfecundity in women but not in men*. Occupational Environmental Medicine 1999; 56:443-448.
- ⁴³ Svensson B-G, Nise G, Erfurth E-M, et al. *Hormone status in occupational toluene exposure*. American Journal of Industrial Medicine 1992; 22:99-107.
- ⁴⁴ Eskenazi B, Fenster L, Hudes M, et al. *A study of the effects of perchloroethylene exposure on the reproductive outcomes of wives of dry-cleaning workers*. American Journal of Industrial Medicine 1991; 20:593-600.
- ⁴⁵ De Cock J, Westveer K, Heederik D, et al. *Time to pregnancy and occupational exposure to pesticides in fruit growers in the Netherlands*. Occupational Environmental Medicine 1994; 51:693-699.
- ⁴⁶ Centers for Disease Control and Prevention Diabetes Public Health Resource. www.cdc.gov/diabetes/statistics/incidence/fig4.htm. Viewed April 9, 2007.
- ⁴⁷ Centers for Disease Control and Prevention National Diabetes Fact Sheet, United States, 2003. www.cdc.gov/diabetes/pubs/pdf/ndfs_2003.pdf. Viewed April 9, 2007.
- ⁴⁸ Newbold RR, Padilla-Banks E, Snyder RJ, Jefferson WN. *Developmental exposure to estrogenic compounds and obesity*. Birth Defects Research. Part A, Clinical Molecular Teratology 2005 Jul; 73(7):478-80.
- ⁴⁹ Tseng CH, Tseng CP, Chiou HY, Hsueh YM, Chong CK, Chen CJ. *Epidemiologic evidence of diabetogenic effect of arsenic*. Toxicology Letters 2002 Jul 7; 133(1):69-76.
- ⁵⁰ Bertazzi PA, Consonni D, et al. *Health effects of dioxin exposure: a 20-year mortality study*. American Journal of Epidemiology 2001 Jun 1; 153(11):1045-7; discussion 1048-9.
- ⁵¹ Rene B. J. Remillard and Nigel J. Bunce. *Linking dioxins to diabetes: epidemiology and biologic plausibility*. Environmental Health Perspectives 2002 Sep; 110(9):853-8.
- ⁵² Alonso-Magdalena, P, S Morimoto, C Ripoll, E Fuentes and A Nadal. *The estrogenic effect of bisphenol-a disrupts the pancreatic β -cell function in vivo and induces insulin resistance*. Environmental Health Perspectives 2006 Jan; 114(1):106-112.
- ⁵³ Lee DH, Lee IK, Jin SH, Steffes M, Jacobs DR Jr. *Association Between Serum Concentrations of Persistent Organic Pollutants and Insulin Resistance Among Nondiabetic Adults: Results from the National Health and Nutrition Examination Survey 1999-2002*. Diabetes Care 2007 Mar; 30(3):622-628.
- ⁵⁴ Mott L. *The disproportionate impact of environmental health threats on children of color*. Environmental Health Perspectives 1995 Sep; 103 Suppl 6:33-5.
- ⁵⁵ National Institute of Health, Office Of Research On Women's Health. *Women of Color Health Data Book. 2006*. <http://orwh.od.nih.gov/pubs/WomenofColor2006.pdf>.
- ⁵⁶ Massart F, Parrino R, Seppia P, Federico G, Saggese G. *How do environmental estrogen disruptors induce precocious puberty?* Minerva Pediatrica 2006 Jun; 58(3):247-54.
- ⁵⁷ Howdeshell KL, Hotchkiss AK, Thayer KA, Vandenberg JG, vom Saal FS. *Exposure to bisphenol A advances puberty*. Nature 1999 Oct 21; 401(6755):763-4.
- ⁵⁸ Parent AS, Rasier G, Gerard A, Heger S, Roth C, Mastronardi C, Jung H, Ojeda SR, Bourguignon JP. *Early onset of puberty: tracking genetic and environmental factors*. Hormone Research 2005; 64 Suppl 2:41-7.
- ⁵⁹ Den Hond E, Schoeters G. *Endocrine disruptors and human puberty*. International Journal of Andrology 2006 Feb; 29(1):264-71; discussion 286-90.
- ⁶⁰ Smith, Allan H., Guillermo Marshall, Yan Yuan, Caterina Ferreccio, Jane Liaw, Ondine von Ehrenstein, Craig Steinmaus, Michael N. Bates, and Steve Selvin. *Increased mortality from lung cancer and bronchiectasis in young adults after exposure to arsenic in utero and in early childhood*. Environmental Health Perspectives 2006 Aug; 114(8):1293-6.
- ⁶¹ Carpenter, David O., Kathleen Arcaro and David Spink. *Understanding the Human Health Effects of Chemical Mixtures*. Environmental Health Perspectives 2002 Feb; 110 Suppl 1:25-42.
- ⁶² Schettler, Ted. *Toward an Ecological View: Complex Systems, Health and Disease*. San Francisco Medicine: Journal of the San Francisco Medical Society February 2006; 79(1).
- ⁶³ Celia Byrne, Ph.D. National Cancer Institute Risk Factors for Breast Cancer Factsheet. http://rex.nci.nih.gov/NCI_Pub_Interface/raterisk/risks120.html. Viewed April 10, 2007.
- ⁶⁴ American Lung Association Air Pollution and Exercise. www.lungusa.org/site/pp.asp?c=dvLUK900E&b=36292. Viewed April 10, 2007.
- ⁶⁵ Campbell ME, Li Q, Gingrich SE, Macfarlane RG, Cheng S. *Should people be physically active outdoors on smog alert days?* Canadian Journal of Public Health 2005 Jan-Feb; 96(1):24-8.
- ⁶⁶ Nelson HD, Huffman LH, Fu R, Harris EL. *Genetic risk assessment and BRCA mutation testing for breast and ovarian cancer susceptibility: systematic evidence review for the U.S. Preventive Services Task Force*. Annals of Internal Medicine 2005 Sep 6; 143(5):362-79.
- ⁶⁷ Ahern MM, Hendryx MS. *Social capital and risk for chronic illnesses*. Chronic Illness 2005 Sep; 1(3):183-90.
- ⁶⁸ Holtgrave DR, Crosby R. *Is social capital a protective factor against obesity and diabetes? Findings from an exploratory study*. Annals of Epidemiology 2006 May; 16(5):406-8.
- ⁶⁹ Drayton-Brooks S, White N. *Health promoting behaviors among African American women with faith-based support*. Association of Black Nursing Faculty Journal 2004 Sep-Oct; 15(5):84-90.
- ⁷⁰ Colborn, Theo. *Neurodevelopment and endocrine disruption*. Environmental Health Perspectives 2004 Jun; 112(9):944-9.
- ⁷¹ Bjornberg KA, Vahter M, Grawe KP, Berglund M. *Methyl mercury exposure in Swedish women with high fish consumption*. The Science of the Total Environment 2005 Apr 1; 341(1-3):45-52.
- ⁷² Mori K, Yoshida K, Hoshikawa S, Ito S, Yoshida M, Satoh M, Watanabe C. *Effects of perinatal exposure to low doses of cadmium or methylmercury on thyroid hormone metabolism in metallothionein-deficient mouse neonates*. Toxicology 2006 Nov 10; 228(1):77-84.
- ⁷³ Debes F, Budtz-Jorgensen E, Weihe P, White RF, Grandjean P. *Impact of prenatal methylmercury exposure on neurobehavioral function at age 14 years*. Neurotoxicology and Teratology 2006 May-Jun; 28(3):363-75.
- ⁷⁴ Stern AH, Jacobson JL, Ryan L, Burke TA. *Do recent data from the Seychelles Islands alter the conclusions of the NRC Report on the toxicological effects of methylmercury?* Environmental Health 2004 Jan 30; 3(1):2.
- ⁷⁵ National Research Council, Committee on the Toxicological Effects of Methylmercury. *Toxicological Effects of Methylmercury*. Washington, DC: National Academy Press, 2000.
- ⁷⁶ Ng SP, Zelikoff JT. *Smoking during pregnancy: Subsequent effects on offspring immune competence and disease vulnerability in later life*. Reproductive Toxicology 2006 Nov 26; [Epub ahead of print].
- ⁷⁷ Pattenden S, Antova T, Neuberger M, Nikiforov B, De Sario M, Grize L, Heinrich J, Hruby F, Janssen N, Luttmann-Gibson H, Privalova L, Rudnai P, Splichalova A, Zlotkowska R, Fletcher T. *Parental smoking and children's respiratory health: independent effects of prenatal and postnatal exposure*. Tobacco Control 2006 Aug; 15(4):294-301.

- ⁷⁸ Gunes T, Koklu E, Yikilmaz A, Ozturk MA, Akcakus M, Kurtoglu S, Coskun A, Koklu S. *Influence of maternal smoking on neonatal aortic intima-media thickness, serum IGF-I and IGFBP-3 levels*. European Journal of Pediatrics 2007 Jan 4; [Epub ahead of print].
- ⁷⁹ Bailey BA, Byrom AR. *Factors Predicting Birth Weight in a Low-Risk Sample: The Role of Modifiable Pregnancy Health Behaviors*. Maternal and Child Health Journal 2007 Mar; 11(2):173-9.
- ⁸⁰ Perera FP, Rauh V, Whyatt RM, Tang D, Tsai WY, Bernert JT, Tu YH, Andrews H, Barr DB, Camann DE, Diaz D, Dietrich J, Reyes A, Kinney PL. *A summary of recent findings on birth outcomes and developmental effects of prenatal ETS, PAH, and pesticide exposures*. Neurotoxicology 2005 Aug; 26(4):573-87.
- ⁸¹ Dugandzic R, Dodds L, Stieb D, Smith-Doiron M. *The association between low level exposures to ambient air pollution and term low birth weight: a retrospective cohort study*. Environmental Health 2006 Feb 17; 5:3.
- ⁸² Marozieni L, Grazuleviciene R. *Maternal exposure to low-level air pollution and pregnancy outcomes: a population-based study*. Environmental Health 2002 Dec 9; 1(1):6.
- ⁸³ Rogers JF, Dunlop AL. *Air pollution and very low birth weight infants: a target population?* Pediatrics 2006 Jul; 118(1):156-64.
- ⁸⁴ Birnbaum LS, Fenton SE. *Cancer and developmental exposure to endocrine disruptors*. Environ Health Perspectives 2003 Apr; 111(4):389-94.
- ⁸⁵ Murray TJ, Maffini MV, Ucci AA, Sonnenschein C, Soto AM. *Induction of mammary gland ductal hyperplasias and carcinoma in situ following fetal bisphenol A exposure*. Reproductive Toxicology 2006 Oct 24.
- ⁸⁶ Vandenberg LN, Maffini MV, Wadia PR, Sonnenschein C, Rubin BS, Soto AM. *Exposure to environmentally relevant doses of the xenoestrogen bisphenol-A alters development of the fetal mouse mammary gland*. Endocrinology 2007 Jan; 148(1):116-27.
- ⁸⁷ Rushton L. *Health impact of environmental tobacco smoke in the home*. Reviews on Environmental Health 2004 Jul-Dec; 19(3-4):291-309.
- ⁸⁸ Rushton L, Courage C, Green E. *Estimation of the impact on children's health of environmental tobacco smoke in England and Wales*. Journal of the Royal Society of Health 2003 Sep; 123(3):175-80.
- ⁸⁹ Hawamdeh A, Kasasbeh FA, Ahmad MA. *Effects of passive smoking on children's health: a review*. Eastern Mediterranean Health Journal 2003 May; 9(3):441-7.
- ⁹⁰ U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta, Georgia; 2006. www.surgeongeneral.gov/library/secondhandsmoke/report/
- ⁹¹ Rice DC. *Developmental lead exposure: neurobehavioral consequences*. In: Handbook of Developmental Neurotoxicology (Slikker W, Chang LW, eds). Academic Press, San Diego, CA; 1998.
- ⁹² Needleman HL, Reiss JA, Tobin MJ, Biesecker G, Greenhouse J. *Bone lead levels and delinquent behavior*. Journal of the American Medical Association 1996; 275:363-369.
- ⁹³ Darnerud P, Eriksen G, Johannesson T, Larsen P, Viluksela M. *Polybrominated diphenyl ethers: occurrence, dietary exposure, and toxicity*. Environmental Health Perspectives 2001; 109(suppl 1):49-68.
- ⁹⁴ Eriksson P, Viberg H, Jakobsson E, Orn U, Fredriksson A. *A brominated flame retardant, 2,2',4,4',5-pentabromodiphenyl ether: uptake, retention, and induction of neurobehavioral alterations in mice during a critical phase of neonatal brain development*. Toxicological Sciences 2002 May; 67(1):98-103.
- ⁹⁵ Grandjean P, Landrigan PJ. *Developmental neurotoxicity of industrial chemicals*. Lancet 2006 Dec 16; 368(9553):2167-78.
- ⁹⁶ National Research Council, Committee on Pesticides in the Diets of Infants and Children, Board on Agriculture and Board on Environmental Studies and Toxicology, Commission on Life Sciences. *Pesticides in the Diets of Infants and Children*. National Academy Press, Washington, D.C. 1993.
- ⁹⁷ Schettler T. *Human exposure to phthalates via consumer products*. International Journal of Andrology 2006 Feb; 29(1):134-9; discussion 181-5.
- ⁹⁸ Koo HJ, Lee BM. *Estimated exposure to phthalates in cosmetics and risk assessment*. Journal of Toxicology and Environmental Health, Part A 2004 Dec; 67(23-24):1901-14.
- ⁹⁹ Duty SM, Ackerman RM, Calafat AM, Hauser R. *Personal care product use predicts urinary concentrations of some phthalate monoesters*. Environmental Health Perspectives 2005 Nov; 113(11):1530-5.
- ¹⁰⁰ American Lung Association Adolescent Smoking Statistics. www.lungusa.org/site/pp.asp?c=dvLUK900E&b=39868. Viewed April 10, 2007.
- ¹⁰¹ Marsit CJ, Kim DH, Liu M, Hinds PW, Wiencke JK, Nelson HH, Kelsey KT. *Hypermethylation of RASSF1A and BLU tumor suppressor genes in non-small cell lung cancer: implications for tobacco smoking during adolescence*. International Journal of Cancer 2005 Mar 20; 114(2):219-23.
- ¹⁰² Miller MD, Marty MA, Broadwin R, Johnson KC, Salmon AG, Winder B, Steinmaus C. *The association between exposure to environmental tobacco smoke and breast cancer: A review by the California Environmental Protection Agency*. Preventive Medicine 2007 Feb; 44(2):93-106.
- ¹⁰³ Clapp, Richard, Genevieve Howe and Molly Jacobs. *Environmental and Occupational Causes of Cancer*. Boston University School of Public Health and Environmental Health Initiative, University of Massachusetts Lowell. 2005.
- ¹⁰⁴ Rier, S. and W.G. Foster, *Environmental dioxins and endometriosis*. Seminars in Reproductive Medicine 2003; 21(2): p. 145-54.
- ¹⁰⁵ Rier S, Turner W, Martin D, et al. *Serum levels of TCDD and dioxin-like chemicals in Rhesus monkeys chronically exposed to dioxin: correlation of increased serum PCB levels with endometriosis*. Toxicological Sciences 2001; 59:147-159.
- ¹⁰⁶ Mayani A, Barel S, Soback S, et al. *Dioxin concentrations in women with endometriosis*. Human Reproduction 1997; 12:373-375.
- ¹⁰⁷ Eskenazi B, Mocarelli P, Warner M, et al. *Serum Dioxin Concentrations and Endometriosis: A Cohort Study in Seveso, Italy*. Environmental Health Perspectives 2002; 110(7):629-634.
- ¹⁰⁸ Birnbaum L. *The mechanism of dioxin toxicity: relationship to risk assessment*. Environmental Health Perspectives 1994; 102(suppl 9):157-167.
- ¹⁰⁹ Birnbaum LS, Cummings A. *Dioxins and endometriosis: a plausible hypothesis*. Environmental Health Perspectives 2002; 110(1):15-21.
- ¹¹⁰ Rier S., Martin D, Bowman R, et al. *Endometriosis in Rhesus monkeys (Macaca mulatta) following chronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin*. Fundamental and Applied Toxicology 1993; 21:433-441.
- ¹¹¹ Fanton J, Golden J. *Radiation-induced endometriosis in Macaca mulatta*. Radiation Research 1991; 126: 141-146.
- ¹¹² Palumbo G, Varriale L, Paba V, et al. *Effect of space radiation on expression of apoptosis-related genes in endometrial cells: a preliminary study*. Physica Medica 2001; 17 Suppl 1:241-6.
- ¹¹³ Gotz F, Thieme S, Dorner G. *Female infertility--effect of perinatal xenoestrogen exposure on reproductive functions in animals and humans*. Folia Histochemica et Cytobiologica 2001; 39 Suppl 2:40-3.
- ¹¹⁴ Foster WG. *Environmental toxicants and human fertility*. Minerva Ginecologica 2003 Oct; 55(5):451-7.
- ¹¹⁵ Younglai EV, Holloway AC, Foster WG. *Environmental and occupational factors affecting fertility and IVF success*. Human Reproduction Update 2005 Jan-Feb; 11(1):43-57.

- ¹¹⁶ Kumar S. *Occupational exposure associated with reproductive dysfunction*. Journal of Occupational Health 2004 Jan; 46(1):1-19.
- ¹¹⁷ Midasch O, Drexler H, Hart N, Beckmann MW, Angerer J. *Transplacental exposure of neonates to perfluorooctanesulfonate and perfluorooctanoate: a pilot study*. International Archives of Occupational and Environmental Health 2007 Jan 12. [Epub ahead of print]
- ¹¹⁸ Environmental Working Group. *Body Burden—The Pollution in Newborns*. 2005. www.ewg.org/reports/bodyburden2/execsumm.php. Viewed April 9, 2007.
- ¹¹⁹ For more reading, see The Collaborative on Health and the Environment's infertility resource page. www.healthandenvironment.org/infertility/val-lombrosa_documents.
- ¹²⁰ John EM, Savitz DA, Shy CM. *Spontaneous abortions among cosmetologists*. Epidemiology 1994 Mar; 5(2):147-55.
- ¹²¹ Doyle P, Roman E, Beral V, Brookes M. *Spontaneous abortion in dry cleaning workers potentially exposed to perchloroethylene*. Occupational and Environmental Medicine 1997 Dec; 54(12):848-53.
- ¹²² Labreche F, Forest J, Trottier M, Lalonde M, Simard R. *Characterization of chemical exposures in hairdressing salons*. Applied Occupational and Environmental Hygiene 2003 Dec; 18(12):1014-21.
- ¹²³ Shanna H, Swan, Katharina M, Main, Fan Liu, Sara L, Stewart, Robin L, Kruse, Antonia M, Calafat, Catherine S, Mao, J, Bruce Redmon, Christine L, Ter-nand, Shannon Sullivan, J, Lynn Teague, and the Study for Future Families Research Team. *Decrease in anogenital distance among male infants with pre-natal phthalate exposure*. Environmental Health Perspectives 2005 Aug; 113(8):1056-61.
- ¹²⁴ Mendelsohn ME. *Protective effects of estrogen on the cardiovascular system*. American Journal of Cardiology 2002 Jun 20; 89(12A):12E-17E; discussion 17E-18E.
- ¹²⁵ Macrae IM, Carswell HV. *Oestrogen and stroke: the potential for harm as well as benefit*. Biochemical Society Transactions 2006 Dec; 34(Pt 6):1362-5.
- ¹²⁶ Rosano GM, Panina G. *Oestrogens and the heart*. Therapie 1999 May-Jun; 54(3):381-5.
- ¹²⁷ Vahter M, Berglund M, Akesson A. *Toxic metals and the menopause*. Journal of the British Menopause Society 2004 Jun; 10(2):60-4.
- ¹²⁸ Silbergeld EK, Flaws JA. *Chemicals and menopause: effects on age at menopause and on health status in the postmenopausal period*. Journal of Women's Health 1999 Mar; 8(2):227-34.
- ¹²⁹ Ginaldi L, De Martinis M, D'Ostilio A, Marini L, Loreto MF, Martorelli V, Quaglino D. *The immune system in the elderly: II. Specific cellular immunity*. Immunological Research 1999; 20(2):109-15.
- ¹³⁰ See the Collaborative on Health and Environment Toxicant and Disease Database for more information: <http://database.healthandenvironment.org/index.cfm>. Enter "Immune Suppression" in the "Browse by Disease" section.
- ¹³¹ Environmental Protection Agency Age Healthier, Breath Easier. www.epa.gov/aging/resources/factsheets/ahbe_english_2004_0330.pdf. Viewed April 10, 2007.
- ¹³² See Environmental Protection Agency Age Healthier, Breath Easier. www.epa.gov/aging/resources/factsheets/ahbe_english_2004_0330.pdf. Viewed April 10, 2007.
- ¹³³ Hayes TB, Collins A, Lee M, Mendoza M, Noriega N, Stuart AA, Vonk A. *Hermaphroditic, demasculinized frogs after exposure to the herbicide atrazine at low ecologically relevant doses*. Proceedings of the National Academy of Sciences of the United States of America 2002 Apr 16; 99(8):5476-80.
- ¹³⁴ Hayes T, Haston K, Tsui M, Hoang A, Haeffele C, Vonk A. *Herbicides: feminization of male frogs in the wild*. Nature 2002 Oct 31; 419(6910):895-6.
- ¹³⁵ Hayes T, Haston K, Tsui M, Hoang A, Haeffele C, Vonk A. *Atrazine-induced hermaphroditism at 0.1 ppb in American leopard frogs (Rana pipiens): laboratory and field evidence*. Environmental Health Perspectives 2003 Apr; 111(4):568-75.
- ¹³⁶ Dunn, Kyla. *Teenie Weenies*. Frontline Online. www.pbs.org/wgbh/pages/frontline/shows/nature/gallery/alligators.html. Viewed April 10, 2007.
- ¹³⁷ Guillette LJ Jr, Gross TS, Masson GR, Matter JM, Percival HF, Woodward AR. *Developmental abnormalities of the gonad and abnormal sex hormone concentrations in juvenile alligators from contaminated and control lakes in Florida*. Environmental Health Perspectives 1994 Aug; 102(8):680-8.
- ¹³⁸ Guillette LJ Jr, Pickford DB, Crain DA, Rooney AA, Percival HF. *Reduction in penis size and plasma testosterone concentrations in juvenile alligators living in a contaminated environment*. General and Comparative Endocrinology 1996 Jan; 101(1):32-42.
- ¹³⁹ Semenza JC, Tolbert PE, Rubin CH, Guillette LJ Jr, Jackson RJ. *Reproductive toxins and alligator abnormalities at Lake Apopka, Florida*. Environmental Health Perspectives 1997 Oct; 105(10):1030-2.
- ¹⁴⁰ Environmental Working Group. *Canaries in the Kitchen*. www.ewg.org/reports/toxicteflon/es.php. Viewed April 10, 2007.
- ¹⁴¹ Environmental Protection Agency Basic Information on PFOA. www.epa.gov/opptintr/pfoa/pubs/pfoainfo.htm#concerns. Viewed April 10, 2007.
- ¹⁴² Sonne C, Leifsson PS, Dietz R, Born EW, Letcher RJ, Hyldstrup L, Riget FF, Kirkegaard M, Muir DC. *Xenoendocrine pollutants may reduce size of sexual organs in East Greenland polar bears (Ursus maritimus)*. Environmental Science and Technology 2006 Sep 15; 40(18):5668-74.

