

SELECTED REFERENCES ON CHEMICAL INTOLERANCE AND TILT
BY CLAUDIA S. MILLER, M.D. M.S.

Book

Ashford NA, **Miller CS**: *Chemical Exposures: Low Levels and High Stakes*. Second edition. New York, John Wiley and Sons, Inc., 1998.

Book Chapters

Miller CS, Ashford NA: Multiple Chemical Intolerance and Indoor Air Quality. In *Indoor Air Quality Handbook*. Spengler J, Samet J and McCarthy J, Eds., New York, McGraw-Hill, Inc., 2001.

Miller CS: Chemical hypersensitivity and multiple chemical hypersensitivity syndrome. In *Handbook of Olfaction and Gestation*, 2nd Edition. Doty R, Ed., New York, Marcel Dekker, Inc., 2000.

Papers

Miller CS: Toxicant-induced loss of tolerance: Mechanisms of action of addictive stimuli. *Addiction* 96(1):115-139, 2000.

Miller CS: Are we on the threshold of a new theory of disease? Toxicant-induced loss of tolerance and its relationship to addiction and abidction. *Toxicology and Industrial Health* 15:284-294, 1999.

Miller CS, Prihoda TJ: The Environmental Exposure and Sensitivity Inventory (EESI): a standardized approach for measuring chemical intolerances for research and clinical applications. *Toxicology and Industrial Health* 15:370-385, 1999.

Miller CS, Prihoda TJ: A controlled comparison of symptoms and chemical intolerances reported by Gulf War veterans, implant recipients and persons with multiple chemical sensitivity. *Toxicology and Industrial Health* 15:386-397, 1999.

Miller CS: Toxicant-induced loss of tolerance: an emerging theory of disease? *Environmental Health Perspectives* 105(2):445-453, 1997.

Miller CS, Ashford NA, Doty R, Lamielle M, Otto D, Rahill A, Wallace L: Empirical approaches for the investigation of toxicant-induced loss of tolerance. *Environmental Health Perspectives* 105(2):515-519, 1997.

ADDITIONAL REFERENCES

Bailer J, Witthoft M, Rist F: The Chemical Odor Sensitivity Scale: reliability and validity of a screening instrument for idiopathic environmental intolerance. *Journal of Psychosomatic Research* 61(1):71-79, 2006.

Bell IR, **Miller CS**, Schwartz GE: An olfactory-limbic model of multiple chemical sensitivity syndrome: possible relationships to kindling and affective spectrum disorders. *Biological Psychiatry* 32(3):218-42, 1992.

Caress SM, Steinemann AC: A national population study of the prevalence of multiple chemical sensitivity. *Archives of Environmental Health* 59(6):300-305, 2004.

Hojo S, Kumano H, Yoshino H, Kakuta K, Ishikawa S: Application of Quick Environment Exposure Sensitivity Inventory (QEESI) for Japanese population: study of reliability and validity of the questionnaire. *Toxicology and Industrial Health* 19(2-6):41-49, 2003.

McKeown-Eyssen G, Baines C, Cole DE, Riley N, Tyndale RF, Marshall L, Jazmaji V. Case-control study of genotypes in multiple chemical sensitivity: CYP2D6, NAT1, NAT2, PON1, PON2 and MTHFR. *International Journal of Epidemiology* 33(5):971-978, 2004.

Overstreet DH, **Miller CS**, Janowsky DS, Russell RW: A potential animal model of multiple chemical sensitivity with cholinergic supersensitivity. *Toxicology* 111:119-134, 1996.

Sorg BA, Willis JR, See RE, Hopkins B, Westberg HH: Repeated low-level formaldehyde exposure produces cross-sensitization to cocaine: possible relevance to chemical sensitivity in humans. *Neuropsychopharmacology* 18(5):385-94, 1998.